Current Status: Active PolicyStat ID: 10208853



 Origination:
 8/7/2018

 Last Approved:
 8/17/2021

 Last Revised:
 8/7/2018

 Next Review:
 8/16/2024

 Owner:
 Erin Eicke: Abilene

ner: Erin Eicke: Abilene
Market Director.

Surgical Services

Policy Area: Surgical Services

Standards & Regulations:

References:

# Traffic Control during Total Joint Replacement Surgery, 4.6669

## **PURPOSE:**

Reducing traffic in the operating room during total joint procedures has proven to be advantageous in the prevention of prosthetic joint infections.

# **POLICY:**

Perioperative personnel will adhere to principles of sterile and aseptic technique during total joint procedures. Principles include but may not be limited to:

- 1. Traffic control and conversations
  - a. Radiology personnel
  - b. Vendor presence
  - c. Poorly timed breaks among OR and anesthesia personnel

#### **Procedure Interventions**

- 1. Referring to preference card prior to opening the supplies for the total joint procedure, ensure all necessary supplies and equipment are in the Operating Room.
- 2. Ensure that x-ray equipment is available, set up and in proper working order before opening the supplies for the case.
- 3. Ensure that the vendor is present and all implantable items or any other vendor provided widget is in the operating room.
- 4. Open the sterile supplies as close to the time of the procedure as possible; ensure that doors are closed; do not abandon the room after supplies are opened. The case is considered started when the sterile supplies are opened.
- 5. Ensure that after the incision is made, door openings are limited to those that are imperative to the case. If personnel need to enter or exit the room, this will be door through the door into/from the sub-sterile corridor.

# References:

Mejia, E., Williams, A., Long, M., (2015). Decreasing prosthetic joint surgical site infections: An interdisciplinary approach. *AORN Journal 101*(2), 213-222. doi: 10.1016/j.aorn.2014.03.017

Panahi, P., et al (2012). Operating room traffic is a major concern during total joint arthroplasty. *Clinical Orthopaedics and Related Research 470*, 2690-2694. doi: 10.1007/s11999-012-2252-4

#### **Attachments**

No Attachments

### **Approval Signatures**

Approver	Date
Tracey Carrigan: Director, Surgical Services	8/17/2021
Heather Hess: Nurse Manager, Acute Nursing	8/3/2021